arXiv:0810.3083v2 [hep-ph] 24 Feb 2009

Phenomenological Implications of a Class of Lepton Mass Matrices

S. Dev^{*,1}, Sanjeev Kumar^{†,2}, Surender Verma^{‡,1} and Shivani Gupta^{§,1}

^{1}*Department of Physics, Himachal Pradesh University, Shimla 171005, INDIA.*
^{2}*School of Physics and Material Science, Thapar University, Patiala 147004, INDIA.*

Abstract

Phenomenological implications of a class of lepton mass matrices with parallel texture structure have been examined and phenomenologically interesting constraints on charged lepton and neutrino mass matrix parameters have been obtained.

1 INTRODUCTION

Mass matrices provide important tools for the investigation of the underlying symmetries and the resulting dynamics. The first step in this direction is the reconstruction of the neutrino mass matrix in the flavor basis. However, the reconstruction results in a large variety of possible structures of mass matrices depending strongly on the mass scale, mass hierarchy and the Majorana phases. In the flavor basis, the mass matrix for Majorana neutrinos contains nine physical parameters viz. three mass eigenvalues, three mixing angles and the three CP-violating phases. The two squared-mass differences (Δm^2_{12} and Δm^2_{13}) and the two mixing angles (θ_{12} and θ_{23}) have been measured in solar, atmospheric and reactor experiments. The third mixing angle θ_{13} and the Dirac-type CP-violating phase δ are expected to be measured in the forthcoming neutrino oscillation experiments. The possible measurement of the effective Majorana mass in neutrinoless double decay searches will provide an additional constraint on the remaining three neutrino parameters viz. the neutrino mass scale and two

---

*dev5703@yahoo.com
†sanjeev.verma@thapar.edu
‡s_7verma@yahoo.co.in
§shiroberts_1980@yahoo.co.in

Majorana CP-violating phases. While the neutrino mass scale will be independently determined by the direct beta decay searches and cosmological observations, the two Majorana phases will not be uniquely determined even if the absolute neutrino mass scale is known. Thus, it is not possible to fully reconstruct the neutrino mass matrix with the observations from the presently feasible experiments. Under the circumstances, it is natural to employ other theoretical inputs for the reconstruction of the neutrino mass matrix. The possible forms of these additional theoretical inputs are constrained by the existing neutrino data. Several proposals have been made in the literature to restrict the form of the neutrino mass matrix by reducing the number of free parameters which include presence of texture zeros [1, 2, 3, 4, 5, 6, 7, 8], requirement of zero determinant [9, 10] and the zero trace condition [11] amongst others. Some attempts aimed at understanding the pattern of quark/lepton masses and mixings by introducing Abelian or non-Abelian flavor symmetries, naturally, lead to texture zeros in the mass matrices. However, the current low energy data are consistent with a limited number of texture zero schemes [1, 2, 3, 4, 5, 6, 7, 8]. In particular, the latest low energy data disfavor [12] all quark mass matrices with five or more texture zeros. Four texture zero Ansätze is specially important since it can successfully describe not only the quark but also the lepton sector including the charged lepton and neutrino masses. Furthermore, not all of the texture zeros in the four texture zero Ansätze can be obtained through weak basis transformations and, therefore, have physical implications. In the present work, we examine a special case of four texture zero Ansätze in which the charged lepton mass matrix and Majorana neutrino mass matrix have the same texture zero structure with zero entries at (1,1) and (1,3) places. This Ansätze has been analyzed earlier [13] under the assumption of factorizable phases in the neutrino mass matrix which, however, is not always possible for a general complex symmetric matrix without unnatural fine tuning of the phases. In the context of the Type II see-saw mechanism for example, the phases of the Dirac neutrino mass matrix and the right handed Majorana neutrino mass matrix are assumed to be fine tuned to achieve the cancellation of phases in most of the analyses [14, 15] (and references therein) reported hitherto. In this sense, the analyses reported so far are incomplete. The condition of factorizable phases in the neutrino mass matrix was relaxed in [16]. However, this analysis was confined to the investigation of neutrino hierarchy and effective Majorana mass. In the present work, we have attempted an exhaustive analysis of a class of four-zero texture lepton mass matrices by examining its full implications for the lepton mass matrices.

All the information about lepton masses and mixings is encoded in the hermitian charged lepton mass matrix M_l and the complex symmetric neutrino mass matrix M_ν. In the present work, we consider a special case of four texture zero Ansätze in which the charged lepton mass matrix and Majorana neutrino mass matrix have the same texture zero structure with zero entries at (1,1) and (1,3) places and are given by

$$M_l = \begin{pmatrix} 0 & A_l & 0 \\ A_l & D_l & B_l \\ 0 & B_l & C_l \end{pmatrix} \tag{1}$$

and

$$M_\nu = \begin{pmatrix} 0 & A_\nu & 0 \\ A_\nu & D_\nu & B_\nu \\ 0 & B_\nu & C_\nu \end{pmatrix} \tag{2}$$

respectively. The hermiticity of M_l requires its diagonal elements D_l and C_l to be real whereas the non-diagonal elements A_l, B_l are, in general, complex with $A_l = |A_l|e^{i\phi_1}$, $B_l = |B_l|e^{i\phi_2}$. In contrast, all the non-vanishing elements of the complex symmetric matrix M_ν are, in general, complex.

The hermitian mass matrix, M_l for the charged lepton, is diagonalized by the unitary transformation

$$M_l = V_l M_l^d V_l^\dagger, \tag{3}$$

where $V_l^\dagger = V_l^{-1}$. The hermitian matrix M_l can, in general, be written as

$$M_l = P_l M_l^r P_l^\dagger \tag{4}$$

where P_l is a unitary diagonal phase matrix, $diag(e^{i\phi_1}, 1, e^{i\phi_2})$, and M_l^r is real matrix which can be diagonalized by a real orthogonal matrix O_l:

$$M_l^r = O_l M_l^d O_l^T \tag{5}$$

where the superscript T denotes transposition and M_l^d=$diag(m_e, -m_\mu, m_\tau)$. From Eqs.(3-5), the unitary matrix V_l is given by

$$V_l = P_l O_l. \tag{6}$$

Using the invariants, $\mathrm{Tr}M_l^r$, $\mathrm{Tr}{M_l^r}^2$ and $\mathrm{Det}M_l^r$, the matrix elements $|A_l|$, $|B_l|$ and C_l can be written in terms of the charged lepton masses and D_l as

$$\begin{aligned} C_l &= m_e - m_\mu + m_\tau - D_l, \\ |A_l| &= \left(\frac{m_e m_\mu m_\tau}{C_l}\right)^{\frac{1}{2}}, \\ |B_l| &= \left[\frac{(m_\tau - m_\mu - D_l)(m_\tau + m_e - D_l)(m_\mu - m_e + D_l)}{C_l}\right]^{\frac{1}{2}}. \end{aligned} \tag{7}$$

Here, D_l should be in the range $(m_e - m_\mu) < D_l < (m_\tau - m_\mu)$ for the elements $|A_l|$ and $|B_l|$ to be real.

Using Eq. (7), the elements of the diagonalizing matrix, O_l can be written in terms of charged lepton masses m_e, m_μ, m_τ and charged lepton mass matrix element D_l, which are given by

$$\left.\begin{array}{r}
O_{11}=\sqrt{\frac{m_\mu m_\tau(m_\tau-m_\mu-D_l)}{(m_e-m_\mu+m_\tau-D_l)(m_e+m_\mu)(m_\tau-m_e)}}\\
O_{12}=\sqrt{\frac{m_\tau m_e(m_\tau+m_e-D_l)}{(m_e-m_\mu+m_\tau-D_l)(m_e+m_\mu)(m_\tau+m_\mu)}}\\
O_{13}=\sqrt{\frac{m_e m_\mu(m_\mu-m_e+D_l)}{(m_e-m_\mu+m_\tau-D_l)(m_\mu+m_\tau)(m_\tau-m_e)}}\\
O_{21}=\sqrt{\frac{m_e(m_\tau-m_\mu-D_l)}{(m_e+m_\mu)(m_\tau-m_e)}}\\
O_{22}=\sqrt{\frac{m_\mu(m_\tau+m_e-D_l)}{(m_\mu+m_\tau)(m_e+m_\mu)}}\\
O_{23}=\sqrt{\frac{m_\tau(m_\mu-m_e+D_l)}{(m_\tau-m_e)(m_\mu+m_\tau)}}\\
O_{31}=\sqrt{\frac{m_e(m_\mu-m_e+D_l)(m_\tau+m_e-D_l)}{(m_e-m_\mu+m_\tau-D_l)(m_\mu+m_e)(m_\tau-m_e)}}\\
O_{32}=\sqrt{\frac{m_\mu(m_\mu-m_e+D_l)(m_\tau-m_\mu-D_l)}{(m_e-m_\mu+m_\tau-D_l)(m_\tau+m_\mu)(m_e+m_\mu)}}\\
O_{33}=\sqrt{\frac{m_\tau(m_\tau-m_\mu-D_l)(m_\tau+m_e-D_l)}{(m_e-m_\mu+m_\tau-D_l)(m_\mu+m_\tau)(m_\tau-m_e)}}
\end{array}\right\} \tag{8}$$

If D_l is known the diagonalizing matrix O_l and the charged lepton mass matrix M_l^r are fully determined since the charged lepton masses are known.

The complex symmetric neutrino mass matrix, M_ν is diagonalized by an orthogonal matrix, V_ν

$$M_\nu = V_\nu M_\nu^{diag} V_\nu^T. \tag{9}$$

The neutrino mixing matrix or Pontecorvo-Maki-Nakagawa-Sakata matrix, U_{PMNS}, is given by

$$U_{PMNS} = V_l^\dagger V_\nu. \tag{10}$$

The neutrino mixing matrix U_{PMNS} consists of three non trivial CP-violating phases: the Dirac phase, δ, and two Majorana phases, α and β, and, three neutrino mixing angles viz. θ_{12}, θ_{23} and θ_{13}. The neutrino mixing matrix can be written as product of two matrices characterizing Dirac and Majorana type CP violation

$$U_{PMNS} = UP \tag{11}$$

where U and P are given by

$$U=\begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23}-c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23}-s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23}-c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23}-s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix}, P=\begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\alpha} & 0 \\ 0 & 0 & e^{i(\beta+\delta)} \end{pmatrix}. \tag{12}$$

Using Eqs. (9) and (10), the neutrino mass matrix M_ν can be written as

$$M_\nu = P_l O_l U_{PMNS} M_\nu^{diag} U_{PMNS}^T O_l^T P_l. \tag{13}$$

The two zero textures in M_ν yield two complex equations viz.

$$m_1 a^2 + m_2 b^2 e^{2i\alpha} + m_3 c^2 e^{2i(\beta+\delta)} = 0 \tag{14}$$

and

$$m_1 ad + m_2 bg e^{2i\alpha} + m_3 ch e^{2i(\beta+\delta)} = 0 \tag{15}$$

where the complex coefficients a, b, c, d, g and h are given by

$$\begin{aligned}
a &= O_{11}U_{e1} + O_{12}U_{m1} + O_{13}U_{t1},\\
b &= O_{11}U_{e2} + O_{12}U_{m2} + O_{13}U_{t2},\\
c &= O_{11}U_{e3} + O_{12}U_{m3} + O_{13}U_{t3},\\
d &= O_{31}U_{e1} + O_{32}U_{m1} + O_{33}U_{t1},\\
g &= O_{31}U_{e2} + O_{32}U_{m2} + O_{33}U_{t2},\\
h &= O_{31}U_{e3} + O_{32}U_{m3} + O_{33}U_{t3}.
\end{aligned} \tag{16}$$

Solving Eqs. (14-15) for the two mass ratios $\left(\frac{m_1}{m_2}\right)$ and $\left(\frac{m_1}{m_3}\right)$, we obtain

$$\begin{aligned}
\tfrac{m_1}{m_2}e^{-2i\alpha} &= \tfrac{b(cg-bh)}{a(ah-cd)},\\
\tfrac{m_1}{m_3}e^{-2i\beta} &= \tfrac{c(bh-cg)}{a(ag-bd)}e^{2i\delta}.
\end{aligned} \tag{17}$$

One can enumerate the number of parameters in Eqs. (17). The nine parameters (three neutrino mixing angles $(\theta_{12}, \theta_{23}, \theta_{13})$, three neutrino mass eigenvalues (m_1, m_2, m_3), two Majorana phases (α, β) and one Dirac-type CP violating phase, δ) come from neutrino sector and four parameters (three charged lepton masses (m_e, m_μ, m_τ) and D_l) come from charged lepton sector, thus, totalling 13 parameters. The three charged lepton masses are [17]

$$m_e = 0.510998910MeV, m_\mu = 105.658367MeV, m_\tau = 1776.84MeV. \tag{18}$$

The experimental constraints on the neutrino parameters at one standard deviation are [18]

$$\begin{aligned}
\Delta m^2_{12} &= 7.67^{+0.22}_{-0.21} \times 10^{-5}eV^2,\\
\Delta m^2_{13} &= (2.37 \pm 0.15) \times 10^{-3}eV^2,\\
\theta_{12} &= 34.5 \pm 1.4,\\
\theta_{23} &= 42.3^{+5.1}_{-3.3}.
\end{aligned} \tag{19}$$

Only an upper bound is known on the mixing angle θ_{13} from the CHOOZ experiment. However, the latest global analysis [19] gives

$$\sin^2\theta_{13} = 0.016^{+0.01}_{-0.01} \tag{20}$$

which is non-zero at 90% C.L.. The parameters δ and D_l are varied uniformly within their full ranges. Thus, we are left with three unknown parameters viz. m_1, α, β.

From Eqs. (17), two mass ratios $(\frac{m_1}{m_2})$ and $(\frac{m_1}{m_3})$ can be written as

$$\frac{m_1}{m_2} = \left|\frac{b(cg-bh)}{a(ah-cd)}\right|, \tag{21}$$

$$\frac{m_1}{m_3} = \left| \frac{c(bh - cg)}{a(ag - bd)} e^{2i\delta} \right|. \tag{22}$$

For the simultaneous existence of two texture zeros at $(1,1)$ and $(1,3)$ positions in M_ν, the two values of m_1 given by

$$m_1 = \left(\frac{m_1}{m_2}\right) \sqrt{\frac{\Delta m_{12}^2}{1 - \left(\frac{m_1}{m_2}\right)^2}} \tag{23}$$

and

$$m_1 = \left(\frac{m_1}{m_3}\right) \sqrt{\frac{\Delta m_{12}^2 + \Delta m_{23}^2}{1 - \left(\frac{m_1}{m_3}\right)^2}}, \tag{24}$$

calculated from the mass ratios $\left(\frac{m_1}{m_2}, \frac{m_1}{m_3}\right)$, respectively, must be identical. This constraint can be used to constrain the unknown variables δ, D_l and the third mixing angle θ_{13} which has not been measured experimentally as yet. We have, also, calculated the effective Majorana mass M_{ee} appearing in the neutrinoless double beta decay for the allowed parameter space

$$M_{ee} = m_1|U_{e1}|^2 + m_2|U_{e2}|^2 + m_3|U_{e3}|^2. \tag{25}$$

The Majorana phases α and β are given by

$$\alpha = -\frac{1}{2} arg \left(\frac{b(cg - bh)}{a(ah - cd)}\right), \tag{26}$$

$$\beta = -\frac{1}{2} arg \left(\frac{c(bh - cg)}{a(ag - bd)} \times e^{2i\delta}\right). \tag{27}$$

Using these relations, the Majorana phases α and β have been calculated for the allowed values of δ, θ_{13} and D_l.

2 Results and Discussion

The δ, θ_{13} and D_l parameter space allowed by the current neutrino oscillation data have been depicted in Fig. 1(a) and 1(b) as two dimensional scatter plots. We see that there are two distinct solutions in the (δ, θ_{13}, D_l) parameter space. The more probable solution corresponds to lower values of D_l and will be called "low D_l" solution, henceforth. The less probable solution corresponds to higher values of D_l and will be referred to as the "high D_l" solution in the following discussion. In Fig. 1(c) and 1(d) $M_{ee} - D_l$ and $\alpha - \beta$ correlation plots have been depicted. It is evident from Fig. 1 that these two solutions of D_l have distinguishing implications, especially, for θ_{13}. One of the characterizing feature of low D_l solution is the existence of lower bound on θ_{13} i.e. $\theta_{13} > 2.8^o$. The 95% C.L. ranges of θ_{13} and D_l for these two solutions have been given in Table 1. Majorana mass term M_{ee} is sharply constrained having values of the order of 10^{-3} for low D_l solution, However, for high D_l a wide range of M_{ee} is allowed. The 95% C.L. ranges of M_{ee} for these two solutions are tabulated in Table 1. Also, it can be inferred from Fig. 1 that Dirac-type CP-violating phase δ (Fig. 1(b)) and Majorana phase β (Fig. 1(d)) remain unconstrained for low as well as high D_l solution. However, Majorana phase α is constrained to the range $60^o - 120^o$ for high D_l. In Fig. 2. we have plotted lightest neutrino mass eigenvalue m_1 with θ_{13} for low (left) and high D_l (right) region. This plot has been reported earlier [16], however, no constraints have been obtained on D_l and θ_{13}. It is noted that the analysis done in [16] corresponds to low D_l solution obtained in the present work. We have obtained an upper bound on m_1 of the order of 10^{-2} for low as well as high D_l solution which is in agreement with that obtained in [16]. Also, we find that m_1 is bounded from below for low and high D_l regions for $\theta_{13} < 8^o$ and $< 10^o$, respectively.

It should be noted that in performing numerical analysis we have used value of θ_{13} obtained from global analysis of neutrino data, given by Eq. (20). However, measurement of θ_{13} is the main goal of the future experiments on neutrino oscillations. The upcoming and proposed experiments in this context include Double CHOOZ [20, 21], Daya Bay [22, 23] and RENO [24]. Double CHOOZ is planned to explore $\sin^2 2\theta_{13}$ down to 0.06 in phase-I (0.03 in the later stages) and will be the first to report any observation on θ_{13}. The Daya Bay experiment has a higher sensitivity and plans to observe $\sin^2 2\theta_{13}$ down to 0.01. In this light it will be useful to relax the constraint on θ_{13} [Eq. (20)] and examine the phenomenological implications of this Ansätz for small θ_{13}. We have considered three representative values of $\sin^2 2\theta_{13}$ viz. 0.01, 0.03 and 0.06 and examined the implications for the CP- violating phases α, β and δ depicted in Fig. 3 and Fig. 4, respectively, for low and high D_L solutions. It can be seen from Fig. 3(a) and Fig. 3(b) that as $\sin^2 2\theta_{13}$ is decreased, the Majorana phases α, β becomes more constrained. Furthermore, Dirac-type CP-violating phase δ is constrained to narrower ranges as $\sin^2 2\theta_{13}$ is lowered which is evident from Fig. 4(a). However, for high D_l solution δ remain unconstrained. There is no solution for low D_l with $\sin^2 2\theta_{13} = 0.01$ in Fig. 3. and Fig. 4 because for low D_l the third mixing angle $\theta_{13} > 3^o$ which is, also, evident from Fig. 1(a).

3 Conclusions

In conclusion, we have considered a class of lepton mass matrices with parallel texture structure in charged lepton and Majorana neutrino mass matrices and obtained interesting constraints on the parameters of lepton mass matrices. An interesting feature of this analysis is the rather strong constraints obtained on θ_{13} and the emergence of two classes of solutions corresponding to 'low D_l' and 'high D_l' which is in contrast to the analysis reported earlier [16]. The implications of these two classes have been examined for the unknown parameters of the neutrino mass matrix like θ_{13}, M_{ee} and CP-violating phases without the assumption of factorizable phases in the neutrino mass matrix. Some of the predictions of this analysis are testable in the forthcoming neutrino experiments.

Acknowledgments

The research work of S. D. is supported by the Board of Research in Nuclear Sciences (BRNS), Department of Atomic Energy, Government of India *vide* Grant No. 2004/ 37/ 23/ BRNS/ 399. S. K., S. G. and S. V. acknowledges the financial support provided by Council for Scientific and Industrial Research (CSIR) and University Grants Commission (UGC), Government of India, respectively.

References

[1] Paul H. Frampton, Sheldon L. Glashow and Danny Marfatia, *Phys. Lett.* **B 536**, 79 (2002).

[2] Bipin R. Desai, D. P. Roy and Alexander R. Vaucher, *Mod. Phys. Lett* **A 18**, 1355 (2003).

[3] Zhi-zhong Xing, *Phys. Lett.* **B 530**, 159 (2002).

[4] Wanlei Guo and Zhi-zhong Xing, *Phys. Rev.* **D 67**, 053002 (2003).

[5] Alexander Merle and Werner Rodejohann, *Phys. Rev.* **D 73**, 073012 (2006).

[6] S. Dev and Sanjeev Kumar, hep-ph/0607048.

[7] S. Dev, Sanjeev Kumar, Surender Verma and Shivani Gupta, *Nucl. Phys.* **B 784**, 103 (2007).

[8] S. Dev, Sanjeev Kumar, Surender Verma and Shivani Gupta, *Phys. Rev.* **D 76**, 013002 (2007).

[9] G. C. Branco, R. Gonzalez Felipe, F. R. Joaquim and T. Yanagida, *Phys. Lett.* **B 562** 265 (2003).

[10] Bhag C. Chauhan, Joao Pulido and Marco Picariello, *Phys. Rev.* **D 73**, 053003 (2006).

[11] Xiao-Gang He and A. Zee, *Phys. Rev.* **D 68**, 037302 (2003), hep-ph/0302201 v2.

[12] Monika Randhawa, V. Bhatnagar, P. S. Gill, M. Gupta, *Phys. Rev.* **D 60**, 051301 (2005).

[13] Koichi Matsuda, Hiroyuki Nishiura, *Phys. Rev.* **D 74**, (2006) 033014.

[14] Gulsheen Ahuja, Sanjeev Kumar, Monika Randhawa, Manmohan Gupta, S. Dev, *Phys. Rev.* **D 76**, 013006 (2007).

[15] Z. z. Xing and H. Zhang, *Phys. Lett.* **B 569**, 30 (2003).

[16] G. C. Branco, D. Emmanuel-Costa, R. González Felipe and H.Serôdio, *JHEP* **0801**, 043 (2008), hep-ph/0711.1613v1.

[17] C. Amsler *et al.*, *Phys. Lett.* **B 667**, 1 (2008).

[18] M. C. Gonzalez-Garcia, Michele Maltoni, *Phys. Rept.* 460 (2008) 1-129, hep-ph/0704.1800v2.

[19] G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo, A. M. Rotunno, arXiv:hep-ph/0806.2649.

[20] http://doublechooz.in2p3.fr/.

[21] F. Ardellier *et al.* (Double Chooz Collaboration), hep-ex/0606025.

[22] http://dayabay.ihep.ac.in/.

[23] Daya Bay Collaboration, hep-ex/0701029.

[24] http://neutrino.snu.ac.kr/RENO/.

Parameter	Low D_l	High D_l
θ_{13}	$5.4^o - 11.2^o$	$3.4^o - 11.2^o$
D_l	$(-85 - 415)$ MeV	$(1555 - 1665)$ MeV
M_{ee}	$(1.0 - 1.5) \times 10^{-3}$ eV	$(0.9 - 18.0) \times 10^{-4}$ eV

Table 1: The ranges of θ_{13}, D_l and M_{ee} at 95% C.L..



Figure 1: The parameter space allowed by the current neutrino oscillation data [18] and θ_{13} from global analysis [19]. The scatter plots of θ_{13}, δ, M_{ee} with D_l and of β with α. Low D_l (high D_l) solution is represented by symbol 'plus' ('cross').



Figure 2: $(m_1 - \theta_{13})$ correlation plots for low D_l (left) and high D_l (right).



Figure 3: Correlation plot of the allowed points in $(\alpha - \beta)$ plane for low D_l (left) and high D_l region (right).



4(a) 4(b)

Figure 4: $(D_l - \delta)$ correlation plot for $\sin^2 2\theta_{13} = 0.01$, $\sin^2 2\theta_{13} = 0.03$ and $\sin^2 2\theta_{13} = 0.06$ in low (left) and high (right) D_l region.